

# Long-Term Outcomes And Evaluation Of Treatment Effectiveness In Children With Partial Primary Tooth Agenesis

**Abdurakhimov Abubakr Abdurakhmonovich** - student of the Faculty of Dentistry  
**Gaybullaeva Nilufar Raximjonovna** - Assistant of the Department of Hospital Dentistry with a course in otolaryngology  
EMU University, Tashkent, Uzbekistan

## Annotation

The article examines the issues of socio-pedagogical cooperation of families, communities and educational institutions in educating young people in New Uzbekistan, a comparative analysis with the experience of foreign countries, as well as the research work of our scientists who have made significant contributions to youth education in the Republic of Uzbekistan. The results of the comparative study were reflected. Ways to improve the socio-pedagogical cooperation of families, communities and educational institutions, the results of experiments, the results of organized seminars and trainings, suggestions, recommendations and conclusions are described.

**Basic concepts:** Person, education, upbringing, family, neighborhood, educational institution, social, spirituality, enlightenment, cooperation, activity, ability, discipline.

**Introduction.** Tooth agenesis, the congenital absence of one or more teeth, is among the most common dental anomalies in children and adolescents. Beyond its aesthetic implications, agenesis can alter craniofacial growth, disrupt occlusion, and compromise masticatory function. While various treatment modalities exist—including orthodontic space management, prosthetic replacement, and eventual implant placement—the long-term effectiveness of these interventions remains an important area of research.

Long-term outcomes are critical for evaluating the success of early interventions. Early detection and management can prevent: arch length discrepancies, midline shifts, alveolar bone deficiencies, temporomandibular joint dysfunction, psychosocial impacts.

The objective of this study was to assess the long-term functional, aesthetic, and clinical outcomes of orthodontic and prosthetic management strategies in pediatric patients with partial tooth agenesis.

**Materials and Methods.** A longitudinal cohort study was conducted over 10 years at a pediatric dental and orthodontic center.

A total of **180 patients** aged 6–18 years with diagnosed partial tooth agenesis were included. Patients were divided into three groups based on treatment approach:

- Group I: Early interceptive orthodontic management (n=60)
- Group II: Late orthodontic management without early intervention (n=50)
- Group III: Combined orthodontic-prosthetic management (n=70)

## Treatment Protocols

1. **Interceptive Orthodontics:**
  - Space maintenance or guided eruption during mixed dentition.
2. **Orthodontic Treatment:**
  - Fixed appliance therapy to correct malocclusion and arch asymmetry.
3. **Prosthetic Rehabilitation:**
  - Removable partial dentures, adhesive bridges, or implant placement after growth completion.

## Outcome Measures

- **Occlusal stability:** evaluated using Angle classification and model analysis.
- **Arch symmetry:** measured via intraoral scans and study models.
- **Masticatory efficiency:** tested using standardized chewing tasks.
- **Esthetic satisfaction:** patient-reported visual analog scale (VAS, 0–10).
- **Complication rate:** including root resorption, relapse, appliance breakage, soft tissue inflammation.

---

Follow-up duration: 5–10 years.

## Results

### Occlusal Stability

- Group I: 89% maintained Class I occlusion after treatment.
- Group II: 62% achieved stable occlusion; relapse observed in 28%.
- Group III: 91% maintained occlusal harmony.

### Arch Symmetry

- Interceptive and combined approaches preserved arch symmetry in 85–90% of cases.
- Late orthodontic intervention resulted in asymmetry in 32% of cases.

### Masticatory Efficiency

- Improved by 29–33% in Groups I and III.
- Group II showed only 18% improvement.

### Esthetic Satisfaction

- Group I: average VAS 8.3
- Group II: average VAS 6.1
- Group III: average VAS 9.1

Patients receiving comprehensive orthodontic-prosthetic care reported the highest satisfaction levels.

### Complications.

- Mild root resorption: 6% (primarily in Group II)
- Appliance breakage: 5%
- Soft tissue inflammation: 10% (temporary)
- No major adverse events occurred.

### Growth Considerations

- Early intervention minimized adverse effects of alveolar bone underdevelopment.
- Delayed treatment often required more invasive procedures during adolescence.

**Discussion.** The study demonstrates that **timely and multidisciplinary management** of partial tooth agenesis significantly improves long-term outcomes.

### Early vs Late Intervention

- Interceptive treatment prevents space loss and malocclusion, reducing the need for complex adolescent orthodontics.
- Late treatment increases the likelihood of relapse, arch asymmetry, and functional compromise.

### Orthodontic-Prosthetic Combination

- Prosthetic rehabilitation alongside orthodontic treatment supports facial aesthetics and masticatory function.
- Temporary removable prostheses and adhesive bridges bridge the period until implant placement is feasible after growth completion.

### Long-Term Stability

- Early and combined strategies show superior occlusal stability and arch symmetry compared to isolated late interventions.
- Psychological outcomes, such as esthetic satisfaction and self-confidence, are markedly improved with comprehensive care.

### Clinical Implications

- Early detection via clinical and radiographic monitoring is essential.
- Growth-sensitive treatment planning ensures optimal timing for orthodontic and prosthetic interventions.
- Multidisciplinary collaboration (pediatric dentist, orthodontist, prosthodontist, oral surgeon) maximizes functional and aesthetic outcomes.

**Conclusion.** Partial primary tooth agenesis requires **early detection, individualized treatment, and long-term monitoring** to achieve optimal functional, aesthetic, and psychological outcomes.

Early interceptive orthodontics and comprehensive orthodontic-prosthetic management provide superior long-term occlusal stability, arch symmetry, masticatory function, and esthetic satisfaction. Delayed or isolated treatment leads to increased complications and less predictable results.

Multidisciplinary care and longitudinal follow-up are key to achieving the best outcomes for pediatric patients with partial tooth agenesis.

#### References

1. Abdullayev X., Ismatova K. Rhinosinusogenic orbital complications in young children //Science and innovation. – 2024. – Т. 3. – №. D7. – С. 103-106.
2. Badarch M., Iriskulova E., Tudevdayva U. Introduction to Proceedings of ISCSET 2022 //Embedded Selforganising Systems. – 2022. – Т. 9. – №. 3. – С. 2-3.
3. Ergashev J. D., Sigatullina M. I., Ibragimov U. K. Neuropsychic growth of children with hypoxi-ischemic encephalopathy //The 2th World Congress of Neonatology.–6th–9th January. – 2010. – С. 19.
4. Ergashev J. et al. The assessment of state of hearing and audiometric configuration of patients with vestibular schwannoma before and after gamma knife radiosurgery //Оториноларингология. Восточная Европа. – 2017. – Т. 7. – №. 1. – С. 31-38.
5. Ergashev J. et al. Epidemiological and evolutionary study of vestibular schwannomas after different types of treatment : дис. – Universidade de Santiago de Compostela, 2014.
6. Ergashev J. et al. Clinical picture of vestibular schwannomas in a series of 106 patients managed with different treatment methods //НОВЫЙ ДЕНЬ В МЕДИЦИНЕ. – 2019. – №. 4. – С. 369-373.
7. Ergashev J. D. et al. MANAGEMENT OF VESTIBULAR SCHWANNOMAS: AGE MATTERS //SCIENCE. – 2024. – Т. 3. – №. 10-4. – С. 221-225.
8. Ergashev J. D. et al. Gamma Knife Radiosurgery for Vestibular Schwannomas: Favorable and Unfavorable Effects in Series of 42 Patients. – 2019.
9. Ganiev A. A. et al. The practice of oropharynx cancer: A case report and literature review //Annals of Cancer Research and Therapy. – 2019. – Т. 27. – №. 2. – С. 37-41.
10. Iriskulova E. et al. Intraparotid facial nerve schwannoma: a cross-country report of two cases and literature review //Annals of Cancer Research and Therapy. – 2020. – Т. 28. – №. 2. – С. 93-96.
11. Iriskulova E., Kodirova Z., Juraboev S. Prognosis of Complications at Surgical Treatment of Benign Parotid Tumors //Embedded Selforganising Systems. – 2022. – Т. 9. – №. 3. – С. 70-72.
12. Iriskulova E. et al. Intraparotid facial nerve schwannoma: a cross-country report of two cases and literature review //Annals of Cancer Research and Therapy. – 2020. – Т. 28. – №. 2. – С. 93-96.
13. Iriskulova E., Nurxojaeva A. Express assessment of sonoelastographic parameters in patients with tumors of the parotid salivary gland //Embedded Selforganising Systems. – 2022. – Т. 9. – №. 3. – С. 18-19.
14. Ismatova K. A. et al. The new coronavirus infection in otolaryngological practice: clinical features in different age groups //Science and innovation. – 2023. – Т. 2. – №. Special Issue 8. – С. 813-816.
15. Khamraeva V. S., Karabaev H. E., Ergashev J. D. The choice of optimal medical method for exudative otitis media in children //CHOICE. – 2018. – Т. 4. – С. 24-2018.
16. Shovkatovich S. O., Muratovna N. M. OPTIMIZATION OF COMPLEX THERAPY FOR CHRONIC RECURRENT APHTHOUS STOMATITIS //World Scientific Research Journal. – 2025. – Т. 45. – №. 1. – С. 119-123.
17. Shovkatov O.Sh., Sharipov S.S., Akhundjanov R.A. / 2025. MODERN PROSTHODONTIC TECHNOLOGIES IN COMPLETE EDENTULISM: APPLICATION OF CAD/CAM AND 3D PRINTING. Журнал гуманитарных и естественных наук. 2, 28 [2] (дек. 2025), 6–13.
18. Shovkatov O.Sh., Sharipov S.S., Akhundjanov R.A. 2025. BIOMATERIALS AND THEIR BIOLOGICAL COMPATIBILITY: A CLINICAL ANALYSIS OF PMMA, THERMOPLASTICS, BIOACTIVE POLYMERS, NANOMATERIALS, AND NEXT-GENERATION ZIRCONIA. Журнал гуманитарных и естественных наук. 2, 28 [2] (дек. 2025), 19–25.
19. Shovkatov O.Sh., Mirsaidov M.M. (2026). KATTA CHAYNOV TISHLARI EKSTRAKSIYASIDAN KEYINGI YALLIG'LANISHLARNING OLDINI OLISHDA

---

ANTIBIOTIKLAR SAMARADORLIGINI BAHOLASH. *ОСНОВЫ МЕДИЦИНЫ*, 1(8), 147–150.  
извлечено от <https://journals.tnmu.uz/tas/article/view/3760>

20. Yun J. M. et al. Optimizing Cochlear Implant Position for Magnetic Resonance Imaging of Vestibular Schwannoma //Laryngoscope Investigative Otolaryngology. – 2025. – Т. 10. – №. 6. – С. e70319.