

Biological Characteristics of Fungal Pathogens of Bulb Flowers and Control Measures

Surayyo Misirova and Ibrohim Qurbanov

Namangan Engineering-Technological Institute, Namangan, Uzbekistan

E-mail: samisirova@mail.ru,

Abstract

The experiments were carried out in floriculture farms located in Namangan region during 2017-2019 years, and laboratory work was carried out at floriculture centers in Namangan region of the Republic of Uzbekistan. 19 species and 5 forms of fungal pathogens were detected in 7 species of bulb flowers grown in the Namangan regions of the Republic of Uzbekistan. Infection of selected bulbous flowers with 27 species of pathogenic fungi was revealed. It was revealed that the selected 7 species of bulb flower plants were infected with the most dangerous diseases, such as 5 of them by *botrytis* and rust diseases; 4 by *Fusarium* wilt; 4 by *Fusarium* rot; 1 by *powdery mildew*; 2 by *sclerotiniosis*; 3 by *penicilliosis*, and by 1 *powdery mildew*, black aphids and false *verticillioses* were infected. In the conditions of the Namangan region of the Republic of Uzbekistan, the fungus *Trichoderma viride* was used as a biological method on 7 types of bulbous flowering plants, and fungicides such as Maxim, Bayleton, Fundazol were used as a chemical method in the fight against identified diseases.

1. Introduction

Currently, around the world, efforts are made to combat more than 1,500 species of pests and microorganisms that cause diseases in decorative flowers. 92% of the microorganisms that cause disease are various fungi. Therefore, the definition of diseases of decorative flowers that cause various fungi, their bioecological properties, distribution rules and improvement of measures to combat them makes it possible to improve the efficiency of floriculture [1-3].

Scientists conducted studies on cultivating flower varieties that are new to the climatic conditions of Uzbekistan, developed measures to combat the types of fungal pathogens and achieve resistance to other factors, and developed agricultural techniques for growing healthy decorative flowers. In this regard, great attention was paid to increasing research on the improvement and introduction into production of methods for creating flower varieties resistant to diseases and pests suitable for soil climatic conditions and methods for the prevention of diseases caused by pathogenic fungi [4-6].

At present, in a number of developed countries of the world, including Holland, England, France, and the USA, effective methods of growing flowering plants, determining their diseases, and controlling them have been put into production. It was determined that species of fungal pathogens such as *Fusarium*, *Botrytis*, *Sphaerotheca*, *Puccinia* are geographically widespread, change their shape depending on climatic conditions. For this reason, in order to improve effective methods of combating fungal pathogens of many flowers planted and cultivated in the Republic of Uzbekistan. Many researches dedicated to their bioecological properties, the laws of distribution [7,8].

The following plants can be attributed to the bulbous cultures of spring flowering: tulips, hyacinths, daffodils, decorative bows, fritillaria, muscarias, cannons, crocuses, blueberries, snowdrops and others. They have approximately the same pests, diseases and symptoms of their manifestations. The exception is fritillaria and bows, which do not eat mice and bypass the raspberries. Onion cultures are known about 30 species of fungal, viral and bacterial diseases and harmful insects. Some are rare, others more often. But there is one general rule: the healthier the bulbs, the better the preparatory work carried out with the ground before planting, and the more favorable the climatic conditions, the less various diseases attack

Research Objectives:

1. Collection of herbarium samples of the infected bulb flower and its affected parts and identification of the types of fungal pathogens.
2. Definition of the analysis of the types of fungal pathogens.

3. Determining the degree of infection of selected plants with the most dangerous types of fungal pathogens;
4. Improvement of control measures against the most common types of fungal pathogens found in bulbous flowers.

2. Research methods.

The studies conducted according to current methods in accordance with standards in this field of science. The method of M.K. Khokhryakov used for collecting herbarium samples of bulbous flowers infected with various fungi pathogens in the field. When isolating and determining the types of fungi from seeds of bulbous flowering plants in laboratory conditions, the methods of N.A. Naumov, A.Ya. Semenov, A.P. Abramov, M.K. Khokhryakov. The determination of fungal species in the cells of infected plants was carried out by definition (identifiers) developed by V.I. Bilay, P.N. Golovin, T.A. Dobrozrakova, M.K. Khokhryakov, N.I. Gaponenko.

The object of the research is bulb flowers, widely distributed in certain regions and highly appreciated by consumers, such as cloves, hyacinth, gladiolus, iris, lily, tulip, daffodil and phlox.

According to the laws of nature, plants, microorganisms, as well as fungi, change their bioecological characteristics. Therefore, the study of fungal species is an ongoing process.

On a global scale, in Italy, the pathogen *Phytophthora cotianae* var. *parasitica*. Scientists A. Francaschini, Serris S., A. Foddai found that soil fungi of the *Gerber jamesonii* hybrid species, including species belonging to the *Fusarium* family, cause the greatest harm.

In the conditions of our republic, scientists have been studying diseases of agricultural cults for many years. Including P.K. Ozolin P.N. Golovin, N.G. Zaprametov, M.A. Karimov, S.S. Ramazanova, B.O. Khasanov and their many students, having studied the biological, environmental, physico-biochemical characteristics of the types of fungi that cause the greatest harm to crops, the laws of their distribution and systematics, based on this, developed measures to combat them and put them into practice. The studies were carried out in the following sequence: Herbarium samples of infected bulbous flower plants and their parts were collected by the method of M.K. Khokhryakov; the selection of fungi from the seeds of bulbous flower plants and the determination of their species by the methods of N.N. Naumova, A.Ya. Semenova, A.P. Abramova, M.K. Khokhryakova; species of fungi in the cells of infected plants were determined using determinants V.I. Bilaya, P. N. Golovina and T. A. Dobrozrakova.

$$P = \Pi_k \cdot 100/N$$

Where, P - the spread of the disease, %; N is the number of plants in the experiment, pcs.; P - total number of infected bulbous plants in the experiment, pcs. In the fight against infected bulbous flower plants, the methods of S. N. Maskovets, I. S. Fedorinchik and H. T. Tillaev were used [8-10]. The experiments were carried out in floriculture farms located in Namangan region during 2017-2019, and laboratory work was carried out at floriculture centers in Namangan region (Fig. 1).

3. Results And Discussion

3.1. Fungal diseases of bulbous flower plants and their systematic location. Table 1 shows the systematic arrangement of species of fungi isolated from bulb flower plants. This table shows 19 species and 5 forms of fungal pathogens isolated from bulbous flower plants grown under conditions in Namangan regions for 3 years and revealed that they belong to 4 classes, 7 subspecies, 7 families and 12 orders.




Fig. 1. Place of study (Namangan region, Republic of Uzbekistan)

The largest group consists of fungal species that are part of the class *Deuteromycytina* consisting of 19 species and 5 forms. Then follows the class *Basidiomycotina* - 4 species; *Ascomycotina* - 3 species. The smallest group is *Mastigomycotina*, consisting of 1 species. The most common types of fungi of the *Fuzarium* family were found in the study, having 4 species and 4 forms. Of these, *F.oxysporum f. sp. gladioli (Mass.) Snyd.et Hans*, *F.oxysporum f. sp. lilii Jmle*, *F.oxysporum f. sp. narcissi, Snyd. Et Hans*, *F.oxysporum f. sp. tulipae Apr.* cause fusarium wilt.

Table 1.

The systematic arrangement of species of fungi isolated from bulb flowers (2019-2022)

Class	Order	Family	Category	Types of fungi, their forms and variations
1	2	3	4	5
<i>Mastigomycotina</i>	<i>Peronosporales</i>	<i>Peronosporaceae</i>	<i>Peronospora</i>	<i>P. lilii</i> . Stenina
<i>Ascomycotina</i>	<i>Erysiphales</i>	<i>Erysiphaceae</i>	<i>Erysiphe</i>	<i>E.cichoriacearum f. phlogis</i> Jacz.
			<i>Sclerotinia</i>	<i>S. gladioli (Mass.) Dray.</i> <i>S. tuliporum</i> Kleb
<i>Basidiomycotina</i>	<i>Ustilaginales</i>	<i>Ustilaginaceae</i>	<i>Urocystis</i>	<i>U. gladioli</i> W.G. Smith.
	<i>Uredinales</i>	<i>Pucciniaceae</i>	<i>Uromyces</i>	<i>U. lilii (Link.) Fuck.</i>
			<i>Puccinia</i>	<i>P. gladioli</i> Cast. <i>P. iridis (DC.) Wallroth</i>

<i>Deuteromycotina</i>	<i>Moniliales</i>	<i>Moniliaceae</i>	<i>Oidiopsis</i>	<i>O. phlogis</i> Golov.
			<i>Penicillium</i>	<i>P. gladioli</i> Mc. Cull. et. Thom.
			<i>Botrytis</i>	<i>B. gladiolorum</i> Timmer.
	<i>B. narcissicola</i> Kleb.			
	<i>B. tulipae</i> (Lib.) Horkins.			
	<i>Acervulales</i>	<i>Tuberculariaceae</i>	<i>Fusarium</i>	<i>F.oxysporum</i> f. sp. <i>gladioli</i> (Mass.) Snyd.et Hans.
				<i>F.oxysporum</i> f. sp. <i>lilii</i> Jmle.
				<i>F.oxysporum</i> f. sp. <i>narcissi</i> Snyd. Et Hans.
				<i>F.oxysporum</i> f. sp. <i>tulipae</i> Apr.
	<i>Pycnidiales</i>	<i>Sphaerioideae</i>	<i>Pyllosticta</i>	<i>P. paeoniae</i> Sacc. Et Speg.
<i>Septoria</i>			<i>S. paeoniae</i> West.	
Total: 4	7	7(11)	12(28)	19 types, 5 forms

And also, in the experiments, the types of fungus pathogens of bulbous flower plants, the degree of infection of bulbous flowers, which plant organs are infected, the interaction of pathogens and plant hosts, the spread of diseases and the harm they cause to the flower industry were determined in experiments (table-2). In particular, 5 species of bulbous flower plants were infected with botrythiosis and rust diseases; *Fusarium* wilt-4; *Fusarium* rot-4; *powdery mildew*-1; *scleratiniosis*-2; *penicilliosis*-3, *powdery mildew*, *black aphids* and *false verticillioses* were infected with only one bulb flower plant. In general, it was revealed that selected for research purposes, 7 species of bulb flowers were infected with 27 species of fungal pathogens. In the process of conducting research, for the first time in the conditions of the Republic of Uzbekistan in the Namangan region, 27 identified diseases were analyzed and the characteristics of the morphological properties of fungal species were determined.

Table 2.
Names of diseases identified in bulbous flower plants (2019-2022)

Bulb Flower Names	Disease names										
	<i>Botirithiosis</i>	<i>Verticilliosis</i>	<i>mildew</i>	<i>Black aphid</i>	<i>Penicilliosis</i>	<i>Scleratiniosis</i>	<i>Downy mildew</i>	<i>powdery mildew</i>	<i>fusarium wilting</i>	<i>fusarium rot</i>	Total
Gladiolus	+		+	+	+				+		5
Lily	+		+			+	+		+	+	6
Tulip	+				+				+	+	4
Hyacinth			+		+	+				+	4
Narcissus	+		+						+		3
Iris	+		+							+	3
Phlox		+						+			2
Total	5	1	5	1	3	2	1	1	4	4	27

3.2. Improving control measures against the most common types of fungal pathogens in bulbous decorative flowers. Numerous data are known that the fungi of the Trichoderma family have high antagonistic properties against fungi that cause pathogens and diseases caused by phytopathogenic microorganisms found in crops. To determine the effectiveness of *T. viride* against the disease, fusarium rot isolated from bulb flower plants, *trichoderma* grown on oats was used with a calculation of 60 and 120 kg/l and pure fungal spores in quantities of 4 and 8 g/l. The highest biological efficiency was 68.6%, with an increase in the consumption rate to 120 kg/ha, more healthy plants were obtained.

In the Namangan region of the Republic of Uzbekistan, fungicides were used as measures for the chemical control of the diseases of ornamental bulbous flower plants, which are included in the list of "Chemicals approved for use by the State Chemical Commission" special attention was paid to determining their economic efficiency. The following fungicides Baraka (1.0-2.0 kg/t), Maxim-2.5% (0.2 and 0.4 l/t), Vitavax 200-75 are mainly used against the disease *Fusarium* rot of ornamental bulbous flower plants. (3.0-4.0 kg) and Topsin-M 70% s.p. (1.0-1.5 kg / t). Fundazol 50% s.p. is used as a reference against root rot disease of many plants. (2.0 kg / t) which gives good results.

4. Conclusion

Based on the studies, the following conclusions can be drawn: 19 species and 5 forms of fungal pathogens were detected in 7 species of bulb flowers grown in the Namangan regions of the Republic of Uzbekistan. Infection of selected bulbous flowers with 27 species of pathogenic fungi was revealed. It was revealed that the selected 7 species of bulb flower plants were infected with the most dangerous diseases, such as *botrythiosis* and rust diseases; 5 species of bulb flower plants were infected; *Fusarium wilt-4*; *Fusarium rot-4*; *powdery mildew-1*; *sclerotiniosis-2*; *penicilliosis-3*, *powdery mildew*, black aphids and false verticillioses were infected with only one bulb flower plant. In the conditions of the Namangan region of the Republic of Uzbekistan, the fungus *Trichoderma viride* was used as a biological method on 7 types of bulbous flowering plants, and fungicides such as Maxim, Bayleton, Fundazol were used as a chemical method in the fight against identified diseases.

References:

- 1) Misirova, S. A. "Systematic types of fungi of allocated and determined types from decorative flowers in conditions region Tashkent." *Agricultural sciences* 6.11 (2015): 1387.
- 2) Misirova, S. A. "Determining of the measure disease control ornamental crops during the growing season in the conditions Tashkent region." *Global Journal of Bio-Sciences and Biotechnology* 5.1 (2016): 119-124.
- 3) Abdumutalovna, Misirova Surayyo, and Sarimsaqova Nilufar Sobirjonovna. "Bioecology of Fungi-Pathogens of Flower Crops and the System to Combat Them." *Agricultural sciences* 7.08 (2016): 539.
- 4) MISIROVA, SA, and NN ERNAZAROVA. "FIGHTING MEASURES THE DISEASE CAUSES A VERY DANGEROUS FUNGAL SPECIES WIDESPREAD IN TASHKENT REGION." *International Journal of Botany and Research (IJBR)* 6 (2016): 5-12.
- 5) Misirova, Surayyo. "Technology of growing orchid flowers from seeds." *E3S Web of Conferences*. Vol. 390. EDP Sciences, 2023.
- 6) MISIROVA, SA. "TECHNOLOGY OF CULTIVATION AND REPRODUCTION OF ORNAMENTAL AND UNIQUE ORCHID FLOWER IN NAMANGAN CONDITIONS." *World Bulletin of Social Sciences* 17 (2022): 156-164.
- 7) Urmonovich, Numonov Otabek. "MANGOSTEEN NUTRITIONAL PRICE AND FUNCTIONAL PROPERTIES." *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ* 14.5 (2023): 3-5.
- 8) Misirova, S. A. "BIOLOGICAL CHARACTERISTICS OF FUNGAL SPECIES THAT CAUSE DISEASES OF ONION FLOWERS AND MEASURES TO COMBAT THEM." (2022).
- 9) Misirova, S., and M. Haydarova. "Flowers from Nederland are Considered to Develop in the Climatic Conditions of Uzbekistan and Are Identified the types of Fungus." *Annals of the Romanian Society for Cell Biology* (2021): 5922-5929.

- 10) Misirova, S. A., et al. "Determination types of fungi-pathogens of ornamental flower crops in conditions region Namangan." *ISJ Theoretical & Applied Science*, 10 (66) (2018): 185-189.
- 11) Misirova, S. A., M. U. Davlatova, and Sh O. Tuhtaboeva. "Biological Characteristics of Fungal Pathogens of Bulb Flowers and Control Measures." *JournalNX*: 207-214.
- 12) Misirova, S., et al. "Growing Dutch tulips in Namangan region." *Bulletin of Agrarian Science of Uzbekistan* 1 (2021).
- 13) Sabirov, Ravshan Z., et al. "Volume-sensitive anion channels mediate osmosensitive glutathione release from rat thymocytes." *PLoS One* 8.1 (2013): e55646.
- 14) Rashidovna, Melanova Nazira, and Numonov Otabek Urmonovich. "Comparative Characteristics of the Leaving of Glutathione From Cells of Different Types." *International Journal on Orange Technologies* 2.10: 79-82.
- 15) Melanova, N. R., M. U. Davlatova, and O. Numanov. "The Effect of Extracellular Glutathione on the Regulation of Thymocyte Volume in Rats under Conditions of Hypoosmotic Stress." *Annals of the Romanian Society for Cell Biology* (2021): 7032-7038.
- 16) Sabirov, R.Z., Kurbannazarova, R.S., Melanova, N.R. and Okada, Y., 2010, January. Swelling-induced release of glutathione from rat thymocytes. In *JOURNAL OF PHYSIOLOGICAL SCIENCES* (Vol. 60, pp. S13-S13). 1-11-11 KUDAN-KITA, CHIYODA-KU, TOKYO, 102-0073, JAPAN: SPRINGER TOKYO.
- 17) Melanova, Nazira Rashidovna, and Sayyora Abduqahharovna Yulchiyeva. "EFFECT OF EXTRACELLUIAR GLUTATHIONE ON COLLOID-OSMOTIC LYSIS OF HUMAN RED BLOOD CELLS." *Scientific and Technical Journal of Namangan Institute of Engineering and Technology* 2.2 (2020): 144-149.
- 18) Choriyeva, Nargiza Mamarajabovna, and Nazira Rashidovna Melanova. "STUDY OF LYSIS OF HUMAN ERYTHROCYTES UPON ADMINISTRATION OF GOSSYPOL, MEGOSIN AND BATRIDEN." *Scientific and Technical Journal of Namangan Institute of Engineering and Technology* 1.9 (2019): 55-58.
- 19) Меланова, Назира Рашидовна. "Сравнительная характеристика выхода глутатиона из различных типов клеток." *Universum: химия и биология* 5 (59) (2019): 9-12.
- 20) Насритдинов, Ахмаджон Абдухамидович, and Хусниддин Тургунбоевич Киргизов. "Агрегат для полосной обработки почвы." *Современные научные исследования и инновации* 12 (2015): 412-416.
- 21) Байбобоев, Набижон Гуломович, et al. "Энергоресурсосберегающий комбинированный агрегат для обработки почвы." *Вестник Рязанского государственного агротехнологического университета им. П.А. Костычева* 3 (23) (2014): 42-44.
- 22) Насритдинов, А. А., and А. В. Рязанов. "Оптимальные условия установки углоснима." *Техника в сельском хозяйстве* 6 (2003): 34-35.
- 23) Насритдинов, Ахмаджон Абдухамидович. "Результаты исследования формы лобовой поверхности стойки чизеля-культиватора." *Universum: технические науки* 1 (58) (2019): 18-20.
- 24) Бойбобоев, Набижон Гуломович, and Ахмаджон Насритдинов. "Теоретические определение перемещение частиц почвы по поверхности углоснима." *Science Time* 6 (18) (2015): 84-89.
- 25) Misirova, S. A., M. U. Davlatova, and Sh O. Tuhtaboeva. "Biological Characteristics of Fungal Pathogens of Bulb Flowers and Control Measures." *JournalNX*: 207-214.
- 26) Шамситдинов, Ф. "Результаты опыта." *Защита и карантин растений* 5 (2003): 27-27.
- 27) Абдуалимов, Ш. Х., and Ф. Р. Шамситдинов. "Влияние применения стимуляторов роста на всхожесть семян, рост, развитие и урожайность хлопчатника в условиях светлых сероземных каменистых почв Наманганской области Республики Узбекистан." *Актуальные проблемы современной науки* 5 (2019): 47-51.
- 28) Абдуалимов, Шухрат Хамадуллаевич, and Фазлиддин Расулович Шамситдинов. "НАМАНГАН ВИЛОЯТИНИНГ ҚИР АДІРЛИ ТОШЛОҚ ЕРЛАРИДА ЯНГИ СТИМУЛЯТОРЛАРНИНГ ҒЎЗА БАҒГ ЮЗАСИ ВА ҲОСИЛДОРЛИГИГА ТАЪСИРИ." *Журнал Биологии и Экологии* 1 (2019).

- 29) Urmonovich N. O. MANGOSTEEN NUTRITIONAL PRICE AND FUNCTIONAL PROPERTIES //ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ. – 2023. – Т. 14. – №. 5. – С. 3-5.
- 30) Rashidovna, Melanova Nazira, and Numonov Otabek Urmonovich. "Comparative Characteristics of the Leaving of Glutathione From Cells of Different Types." *International Journal on Orange Technologies* 2.10: 79-82.
- 31) Abdukhamidovich, Nasritdinov Akhmadzhon, and Numonov Otagak Urmonovich. "The Results of Theoretical Studies of the Chisel Cultivator Rack Frontal Surface Shape." *Annals of the Romanian Society for Cell Biology* (2021): 5930-5938.
- 32) Abdukhamidovich, Nasritdinov Akhmadzhon, Muhabbat Davlatova Urmanovna, and Numonov Otabek Urmonovich. "Strip Till Age of Soil for Deuteric Sowing (Second Crop)." *International Journal on Orange Technologies* 3.4 (2021): 71-74.
- 33) Abduhamidovich N. A. et al. MANGOSTIN DARAXTI VA MEVASINI TIBBIYOTDA FOYDALANISH //Journal of new century innovations. – 2023. – Т. 28. – №. 2. – С. 12-14.
- 34) Melanova, N. R., M. U. Davlatova, and O. Numanov. "The Effect of Extracellular Glutathione on the Regulation of Thymocyte Volume in Rats under Conditions of Hypoosmotic Stress." *Annals of the Romanian Society for Cell Biology* (2021): 7032-7038.
- 35) Рахманов, Дилшод Ортикбаевич, and Муҳаббат Ўрмоновна Давлатова. "ОРГАНОЛЕПТИЧЕСКИХ ОЦЕНКА СВОЙСТВ РЫБ И РЫБНЫХ ПРОДУКТОВ." *O'ZBEKISTONDA FANLARARO INNOVATSIYALAR VA ILMIY TADQIQOTLAR JURNALI* 2.14 (2022): 583-585.
- 36) Юсупова, Махпуза Нумановна. "БИОЛОГИЧЕСКИЙ МЕТОД ЗАЩИТЫ РАСТЕНИЙ." *Scientific Impulse* 1.9 (2023): 1460-1464.
- 37) Юсупова, Махпуза Нумановна. "АНОРНИ ЗАРАКУНАНДАЛАРДАН ҲИМОЯЛАШ." *PEDAGOG* 6.4 (2023): 562-567.
- 38) Юсупова, Махпуза Нумановна. "САБЗАВОТ ЭКИНЛАРИГА БИОЛОГИК КУРАШ ҲАҚИДА МУЛОХАЗАЛАР." *Scientific Impulse* 1.9 (2023): 1469-1473.
- 39) Юсупова, Махпуза Нумановна. "ФАРҒОНА ВОДИЙСИ ШАРОИТИДА ИГНА БАРГЛИ ДАРАХТЛАРНИ ЗАРАКУНАНДАЛАРДАН ҲИМОЯЛАШ." *SO 'NGI ILMIY TADQIQOTLAR NAZARIYASI* 6.4 (2023): 316-320.
- 40) O'rmonovna, Davlatova Muhabbat. "MANGOSTIN DARAXTI VA UNING KIMYOVIY XUSUSIYATLARI." *INNOVATION IN THE MODERN EDUCATION SYSTEM* 3.25 (2022): 1-4.
- 41) Юсупова, Махпуза Нумановна. "УФТ: 635 САБЗАВОТ ЭКИНЛАРИГА БИОЛОГИК КУРАШ ҲАҚИДА МУЛОХАЗАЛАР." *Научный импульс*: 355.
- 42) Qurbonov, I. "Tulip varieties imported from the netherlands technology of cultivation of namangan region. galaxy international interdisciplinary research journal (giirj) issn (E): 2347-6915 Vol. 9." (2021).
- 43) Kurbanov, I. G. "CARE OF TULIP VARIETIES OF THE NETHERLANDS IN THE CLIMATIC CONDITIONS OF THE NAMANGAN REGION." *American Journal of Interdisciplinary Research and Development* 6 (2022): 117-120.
- 44) Qurbonov I. E-RECRUITMENT: SOCIAL MEDIA AND RECRUITING //InterConf. – 2021.
- 45) Qurbonov, Ibragim Sharifjonovich. "CLONELY MICRO-CULTIVATION OF PLANTS AND ITS APPLICATION TO AGRICULTURE." *Scientific and Technical Journal of Namangan Institute of Engineering and Technology* 1.4 (2019): 74-78.
- 46) Юсупова М. Н., Ахмедова М. М. МЕВАЛИ ДАРАХТЛАРНИ ЗАРАКУНАНДАЛАРИГА УЙҒУНЛАШГАН КУРАШ ЧОРАЛАРИ //ЖУРНАЛ АГРО ПРОЦЕССИНГ. – 2020. – Т. 2. – №. 8.
- 47) Ходжаев, Ш. Т., Сагдуллаев, А. У., Исаев, О. Б., & Юсупова, М. Н. (2011). Проблемы защиты растений в Узбекистане. *Защита и карантин растений*, (8), 23-24. Юсупова М. Особенности защиты хлопчатника посеянного под пленки от вредных организмов //Автореф. канд. дисс./М. Юсупова–Ташкент. – 2001.

- 48) Ходжаев, Ш. Т., Юсупова, М. Н., Курязов, Ш., & Саттаров, Н. (2008). Перспективы биологической защиты хлопчатника от хлопковой совки. *Сб. трудов.-Ташкент: Таллин*, 44-49.
- 49) Yusupova M. N., Nosirov B. Z. Pests of cotton and straw control at collection //EPRA International Journal of Multidisciplinary Research (IJMR)-Peer Reviewed Journal. – 2020. – Т. 6. – №. 12. – С. 57-61.
- 50) Yusupova M. N., Axmedova M. M. Mevali daraxtlarni zararkunandalariga uygunlashgan kurash choralari //Jurnal JURNAL AGRO PROTSESSING. Data publikatsii. – 2020. – №. 8. – С. 12.
- 51) Yusupova M. N. Biological method of crop protection in the fergana valley //Agrarian science. – 2018. – №. 6. – С. 68-70.
- 52) Urmonovich, Numonov Otabek. "MANGOSTEEN NUTRITIONAL PRICE AND FUNCTIONAL PROPERTIES." *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ* 14.5 (2023): 3-5.
- 53) MN, Yusupova, and B. Z. Nosirov. "Control Of Cotton Pests On Stubble Lands." *International Journal of Applied* 10.2 (2015): 99-108.
- 54) Юсупова М. Н., Тургунова А. Н., Очилов С. Н. Система интегрированной защиты растений //Российский электронный научный журнал. – 2015. – №. 1. – С. 169-174.
- 55) Alimzhanova Z. I., Kadyrova D. S., Yusupova M. N. Ceramic pigments based on raw materials from Uzbekistan //Glass and Ceramics. – 2014. – Т. 70. – №. 11-12. – С. 441-443.
- 56) Yusupova M. N., Gapparov A. M. Biological Method Of Plant Protection In Uzbekistan //The American Journal of Agriculture and Biomedical Engineering. – 2020. – Т. 2. – №. 11. – С. 29-32.
- 57) Rashidovna M. N., Urmonovich N. O. Comparative Characteristics of the Leaving of Glutathione From Cells of Different Types //International Journal on Orange Technologies. – Т. 2. – №. 10. – С. 79-82.
- 58) Юсупова М. Н., Носиров Б. З. БИОЛОГИЧЕСКИЙ МЕТОД ЗАЩИТЫ РАСТЕНИЙ В УЗБЕКИСТАНЕ //Научно-практические пути повышения экологической устойчивости и социально-экономическое обеспечение сельскохозяйственного производства. – 2017. – С. 498-501.
- 59) Urmonovich, N. O. (2023). MANGOSTEEN NUTRITIONAL PRICE AND FUNCTIONAL PROPERTIES. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 14(5), 3-5.
- 60) Yusupova M. et al. Protection of after harvest cultures-as a reservetors of cotton pests //Agriculture and Biology Journal of North America. – 2013. – Т. 4. – №. 5. – С. 576-582.
- 61) Ходжаев, Ш. Т., Юсупова, М. Н., Юлдашев, Ф., Исаев, О. Б., & Шокирова, Г. (2011). Борьба с вредителями хлопчатника на пожнивных культурах в севообороте. *Вестник защиты растений*, (2), 46-52.
- 62) Yusupova M. N. et al. Possibilities of the biological method of cotton plant protection //Agriculture and Biology Journal of North America. – 2011. – Т. 2. – №. 5. – С. 742-744.
- 63) Ходжаев, Ш. Т., Юсупова, М. Н., Юлдашев, Ф., & Жамалов, А. Г. (2010). Хлопковая совка на пожнивных культурах. *Защита и карантин растений*, (12), 22-23.
- 64) Хайдарова, Х. А., Юсупова, М. Н., Ихтиярова, Г. А., & Хайдаров, А. А. ПОЛУЧЕНИЕ ХИТОЗАНА ИЗ ПОДМОРА ПЧЕЛ APIS MILLIFERA. *Сучасний рух науки: тези доп. XI міжнародної науково-практичної інтернет-конференції, 8-9 жовтня 2020 р.-Дніпро, 2020.-Т. 2.-426 с.*, 352.
- 65) Yusupova M., Turgunova A., Ochilov S. INTERGRATED PLANT PROTECTION SYSTEMS.