

# Influence Of Sowing Date And Norms On The Place Of The Lower First Pod Of Lentil Varieties

A A Kurbanov

Researcher, Tashkent State Agrarian University

**Abstract:** This article provides data on the norms and date of sowing, the location of the first pod of lentil varieties. In lentil varieties, the location of the lower first pod was found to be 1.9-3.2 cm higher in the Oltin don variety and 3.0-3.5 cm higher in the Darmon variety when the planting norms were exceeded. When lentil varieties were planted in the autumn and spring, the location of the lower first pod was found to be 0.1–0.2 cm, or homogeneous, and it was found that lentil varieties could be harvested without destroying the crop grown when the sowing norm increased.

**Keywords:** sowing date, sowing norm, “Oldin don”, “Darmon”, location of the first pod.

## Introduction

The standard of living of the country's population is assessed primarily by the fact that the members of society are provided with flour, bread, bakery products, pasta, cereals and confectionery from cereals. An alternative solution to this problem is the correct selection of crops based on their biological and economic potential in the cultivation of agricultural crops and the achievement of ecologically clean and high quality products, enriching the plant gene pool at the expense of new and forgotten agricultural crops with valuable characteristics. As a result of expanding the scope of creation of new varieties of crops, we have access to our own food products, which still replace the products entering our markets through imports.

According to the data, legumes differ from other groups of crops in their ability to fix atmospheric nitrogen and assimilate difficult-to-assimilate phosphorus compounds in the soil. Legumes absorb about 2/3 of the total nitrogen from the atmosphere and 1/3 from the soil when all living conditions are normal. Because legumes have the ability to fix atmospheric nitrogen, they are in great need of more phosphorus and potassium fertilizers [5].

Lentils are an annual herbaceous legume, the height of the stem is 15-75 cm. In terms of biological properties, spring, frost-resistant, shelf life 75-115 days, self-pollinating, used in food and fodder production. Green grass contains 8-10% protein, yield 15-25ts / ha [3; B.18].

Lentil's root 1 m, stem 25-50 cm. The grains are round, flat, yellow, green, pink, gray, brown and black. Lentils are a long-lived plant, the seeds germinate at 3-4 OS, germinate quickly at 7-10 OS. Grows well in porous, loamy and gray soils, does not grow well in heavy, muddy and sandy soils [2; B.10-15].

According to S.E. Traubenberg, lentils are planted with 60-120 kg of seed, sowing depth is 4-6 cm during the period of mowing. After the lawn is dry, anti-plowing works are carried out. When the beans reach 50%, the crop is harvested, dried in a threshing floor and ground in combine harvesters, cleaned. Can be stored when the humidity reaches 14-15% [4; B.62-65].

According to H.N.Atabaeva, J.B.Khudaykulov, lentil varieties were tested in Uzbekistan in 1949 on dry lands. This work lasted 2-4 years. The stem of the “local” variety is 25-28 cm, the 1<sup>st</sup> pod is 13-25 cm, 1000 grains weigh 25-38 g, the shelf life is 75-100 days, the number of pods is 10-20, the seeds are small, the yield is 21.3 c / ha [1; B.212].

## Methods And Materials

Field experiments were conducted on the basis of methodological manuals "Methods of State sorting of agricultural culture" (1989), "Methods of conducting field experiments" (UzPITI, 2007). Statistical analysis of the obtained data was carried out using the program Microsoft Excel and on the basis of BD Dospekhov's manual "Methodology of field experience" (1985) by the method of analysis of variance.

**Sowing dates and rates.** Lentils are planted in autumn (late October, early November), spring (March) and summer (June) in areas free from repeated cereal crops (wheat, barley). Lentils are planted in seedbeds sowing cereals and alfalfa. 2, 3, 4 million seeds were used per hectare, sowing depth was 4-6 cm. Taking into account the local conditions, the sowing rate in the experiments was 2 million units per hectare in the 1<sup>st</sup> variant, 3 million units in the 2<sup>nd</sup> variant, and 4 million units in the 3<sup>rd</sup> variant.

Crops were irrigated during the application period and fed with mineral fertilizers. Harvested when lentils are 60% ripe.

### Results And Discussion

The ability to harvest legumes without destroying them also depends on the location of the lower first leg of the crop, as legumes are branched and the branches may be at the bottom of the stem. On the side branches a lot of pods and grains are formed, the harvesters are not able to harvest on the very low branches, the crop remains in the field.

Experiments analyzed the location of the first lower leg in 2011 and found that when lentil varieties were planted in the fall, the Oltin don When 2 million seeds were sown, the lower first pod was at 18.6 cm and the Darmon variety at 19.1 cm. When sowing norms are increased, the first lower pod is 1.9-3.8 cm high in Oltin don variety and 3.0-3.8 cm high in Darmon variety. Experiments In 2012, when the Oltin don variety was planted in the fall, the location of the lower first leg was 20.8 cm, while the Darmon variety was 20.5 cm. When sowing norms are increased, the first lower pod is 2.1-2.5 cm higher in Oltin don variety and 3.3-3.8 cm higher in Darmon variety.

The experiments analyzed the location of the first lower legume in 2013. When the lentil varieties were planted in the fall, the Oltin don variety was planted at 2.5 seeds, the lower first legume was at 19.5 cm, and the Darmon variety was at 19.8 cm. When sowing norms are exceeded, the first lower pod is 2.0-3.3 cm high in Oltin don variety and 2.7-3.5 cm high in Darmon variety.

**Table 1**

**Influence of sowing dates and norms on the height of placement of the first pod in lentil varieties, cm**

№	Options		Years			Average
	Cultivars	Sowing norms, mln, pc/ha	2011	2012	2013	
<b>Sowing in autumn</b>						
1	Oltin don	2	18,6	20,8	19,5	19,7
2	Oltin don	3	20,5	22,9	21,5	21,6
3	Oltin don	4	22,4	23,3	22,8	22,9
4	Darmon	2	19,1	20,5	19,8	19,8
5	Darmon	3	22,1	23,8	22,5	22,8
6	Darmon	4	22,9	23,8	23,3	23,3
<b>Sowing in spring</b>						
1	Oltin don	2	18,7	21,0	19,7	19,8
2	Oltin don	3	20,7	23,1	21,7	21,8
3	Oltin don	4	22,6	23,6	23,1	23,1
4	Darmon	2	19,3	20,7	19,9	20,0
5	Darmon	3	22,3	24,0	22,7	23,0
6	Darmon	4	23,1	24,0	23,5	23,5

When lentil varieties were planted in spring Experiments analyzed the location of the first lower leg in 2011. When the Oltin don variety sowed 2 million seeds, the lower first leg was located at 18.7 cm and the Darmon variety at 19.3 cm. When sowing norms are exceeded, the first lower pod is 2.0-3.9 cm high in Oltin don variety and 3.0-3.8 cm high in Darmon variety.

Experiments In 2012, when the Oltin don variety was planted in the spring, the location of the first lower bean was 21.0 cm in the Oltin don variety and 20.7 cm in the Darmon variety. The variety is located at a height of 3.3–3.3 cm.

Experiments In 2013, when the Oltin don variety was planted in the spring, the location of the lower first leg was 19.7 cm in the Oltin don variety and 19.9 cm in the Darmon variety. When sowing norms are increased, the first lower pod is 2.0-3.4 cm higher in Oltin don variety and 3.3-3.3 cm higher in Darmon variety. An analysis of the average 3-year data revealed that when both varieties were planted in the fall and spring, the location of the first pod was moderately high (Table 1).

### Conclusion

In lentil varieties, the location of the lower first leg is 1.9-3.2 cm higher in the Oltin don variety and 3.0-3.5 cm higher in the Darmon variety when planting norms are exceeded. When lentil varieties were planted in autumn and spring, the location of the lower first leg was different from 0.1-0.2 cm, or found to be homogeneous, and lentil varieties were found to be harvested without destroying the crop grown when planting rates increased.

### References

1. Atabaeva H.N., Khudoykulov J.B.-Botany, Science and Technology, 2018, P.212
2. Возделывание зернобобовых и крупяных культур в Акмолинской области (Рекомендация). Акмолинск, ТОО «Заречный», 2008, 26 с.
3. Технология весеннего сева зерновых, зернобобовых и масличных культур: [//www.agrosektor.kz/recommendations-scientists](http://www.agrosektor.kz/recommendations-scientists). 2010, p.18.
4. Траубенберг С.Е. Ферментативный гидролиз как инструмент для повышения пищевой ценности продуктов растениеводства. // Хранение и переработка сельхозсырья. 2005. №7. С. 62-65.
5. “Чечевица, семена и посев, сорта -//<http://chudo-ogorod.ru>, 2010.